

Manufacture Date:
Expiration Date:
Charge number:
License number and date: 02.10.2020/11703

PLANT TO BE USED AND HARMFUL ORGANISMS:

Plant Name	Name of harmful organism	Application dosage	Time between the last application and harvest
Wheat	Yellow rust (<i>Puccinia striiformis</i>)	125 ml/da	42 days
Sugar beet	Leaf spot (<i>Cercospora beticola</i>)	125 ml/da	42 days
Chickpea	Chickpea anthracnose Anthracnose (<i>Ascochyta rabiei</i>)	125 ml/da	28 days

PREPARATION FOR THE APPLICATION OF THE PLANT PROTECTION PRODUCT:

After the calculation of the plant protection product according to the recommended dosage, mix it with some water in a separate container. Fill the sprayer's tank half with water. Add the mixture to the tank while the mixer of the machine is working. Continue mixing while filling the tank full with water. The mixing process needs to continue until the application is completed. Use the prepared mixture on the same day.

CALIBRATION:

Before the application, the spraying machine should be calibrated. The amount of mixture to be given to the specific area should be well adjusted in order to provide a good coating in applications. Applications should be carried out in cool hours of the day, under non-windy or slightly windy weather conditions.

CLEANING OF THE SPRAYING MACHINE:

Right after the application is completed, empty the tank of the spraying machine safely. After filling the tank with clean water, run the mixer and spray system to ensure that all parts are washed. Do not perform the washing process near water sources. Do not discharge the washing water and waste into water sources.

USE OF THE PLANT PROTECTION PRODUCT:

Wheat yellow rust: Applications should be started when symptoms of disease are seen. Green parts application is carried out against rust factors. The application should be done in a way that the surface of the leaves and stalk is covered with medicated water. If the conditions are suitable for the development of the disease, a second application can be made, taking into account the duration of action of the product.

Leaf spot disease in sugar beet: Applications are started as soon as the weather conditions are suitable for the disease and the first signs of disease are seen on the leaves in the field. Application is continued at 10-12 days intervals, taking into account the severity of the disease, climatic conditions and the effectiveness of the plant protection product.

Chickpea ascochyta blight fungus: Although it is necessary to start the application if the daily average temperature is 10 °C and the proportional humidity is at least 80%, it should be started practically when the first signs of the disease are seen in the region. Application is continued at 10 days intervals, taking into account the severity of the disease, climatic conditions and the effectiveness of the plant protection product. If there was heavy rain on the day of application, the application should be repeated.

INFORMATION ON RESISTANCE:

The plant protection product named **LATİS®** is a fungicide classified as Group C3,11 + G1,3 according to its mechanism of action. Repeated applications of the plant protection products with the same effect mechanism encourage the development of resistance. For this reason, do not exceed the recommended total application number of **LATİS®** in the same production season in order to delay resistance development. In cases where the application needs to be repeated, pay attention to the use of plant protection products with different effect mechanisms (except Groups C3,11 and G1:3)

MIXABILITY:

It is recommended to make pre-mix trials before mixing large amounts for application.



HEKTAŞ TİCARET T.A.Ş

LATİS®

Suspo-Emulsion (SE)

Group	C3,11	G1,3	Fungicide
-------	-------	------	-----------

100 g/l Azoxystrobin + 80 g/l Tetraconazole

READ THE LABEL FIRST

DO NOT USE AT HOME

KEEP AWAY FROM CHILDREN, FOOD AND ANIMAL FEED.

DO NOT SMOKE, EAT OR DRINK ANYTHING DURING APPLICATION OF THE PLANT PROTECTION PRODUCT.

DO NOT LET HUMAN AND ANIMALS IN THE AREA FOR 1 DAY WHERE THE PLANT PROTECTION PRODUCT IS APPLIED.

USE IT IN THE RECOMMENDED MANNER AND DOSAGE TO AVOID RISKS TO HUMAN HEALTH AND THE ENVIRONMENT.

IT IS STRICTLY FORBIDDEN TO USE OTHER THAN THE RECOMMENDED PRODUCTS.

SHAKE AND RINSE THE USED UP PLANT PROTECTION PRODUCT BOTTLE BY FILLING ¼ OF THE EMPTY PACKAGING WITH CLEAN WATER. DISCHARGE THE RINSING WATER INTO THE SPRAYER TANK. REPEAT THIS PROCESS 3 TIMES.

INTOXICATION SYMPTOMS:

General symptoms of intoxication can be seen.

FIRST AID MEASURES:

Eyes: In case of eye contact, rinse with plenty of water for 15–20 minutes with eyes open. Consult a doctor.

Skin: In case of skin contact, take off the clothing. Wash the parts of the body that have come into contact with the plant protection product with plenty of soap and water for 15–20 minutes. Consult a doctor.

Inhalation: Affected person should be brought to fresh air. If necessary, apply artificial respiration. Consult a doctor.

Ingestion: Rinse mouth with plenty of water and consult a doctor. Unless told by the doctor, the affected person should not be made vomiting. An unconscious patient is never given anything orally and is not allowed to vomit.

ANTIDOTE AND TREATMENT:

There is no specific antidote. Treat symptomatically.

A registered trademark of ® HEKTAŞ TİCARET T.A.Ş.

UZEM (National Poison Center) Tel: 114

Net Amount:
Maximum retail price: Imprint

CONTENTS:
Azoxystrobin
Tetraconazole

HAZARD SIGNS:



ATTENTION

HAZARD STATEMENTS (H):

H410 Very toxic to aquatic life with long lasting effects.

PRECAUTIONARY STATEMENTS (P):

P261 Avoid breathing dust and fumes.

P273 Avoid release to the environment.

P280 Wear protective gloves/protective clothing/eye protection/face protection.

P391 Collect spillage.

P501 Dispose of contents / container in accordance with applicable regulations.

EUH401 To avoid risks to human health and the environment, comply with the instructions for use.

POINTS TO BE CONSIDERED WHILE USING AND STORING:

- It is toxic to bees. Do not use in the blooming season.
- It is toxic to fishes. Avoid contaminating underground and surface waters.
- Keep the plant protection product closed in its package.
- It should only be used for the recommended purposes and doses.

CAUTION: Shake well before use the plant protection product!

STORAGE OF THE PRODUCT:

When stored unopened in the original packaging under normal (cool and dry) conditions, the physical, chemical and biological properties of the product do not change except for the tolerance limits for 2 years.

COMPANY STATEMENT:

Manufacturers guarantee the quality of their products provided that they are sold in original packaging. The manufacturers do not accept any liability for any damage that may occur as a result of improper storage or implementation of their products or failure to comply with the recommendations.

LICENSE OWNER AND MANUFACTURER:

HEKTAŞ TİCARET T.A.Ş
Gebze OSB Mahallesi 700. Sk. No: 711/1
41400 Gebze/KOCAELİ
Tel: (0 262) 751 14 12
Fax: (0 262) 751 14 22

